



Callus Ointment Salicyl-Vaseline 10 %

Active ingredient: Salicyl acid
To be used by adults and children over 12 years

Information leaflet: Information for the user

Read the entire information leaflet carefully before starting to use this medicinal product; it contains important information for you.

Always use this medicinal product exactly as described in this leaflet or exactly as prescribed by your doctor or pharmacist or your medical specialist.

- Keep this information leaflet. You may want to reread it at a later date.
- Ask your pharmacist if you need any additional information or advice.
- Please inform your doctor or pharmacist or medical specialist if you notice side effects. This also applies to side effects not stated in this leaflet. See section 4.
- Please consult a doctor if, after 7 days, you do not feel better or even feel worse.

Information in this leaflet:

1. What is Callus Ointment Salicylic Vaseline 10 % and what is it used for?
2. What must I be aware of before using Callus Ointment Salicylic Vaseline 10 %?
3. How should Callus Ointment Salicylic Vaseline 10 % be used?
4. What are the possible side effects?
5. How should Callus Ointment Salicylic Vaseline 10 % be stored?
6. Pack contents and additional information

1. What is Callus Ointment Salicylic Vaseline 10 % and what is it used for?

Areas of application: Increased callus formation (hyperkeratosis), corns (clavus), calluses (callus).

2. What must I be aware of before using Callus Ointment Salicylic Vaseline 10 %?

Callus Ointment Salicylic Vaseline 10 % must not be used,

- If you are allergic to salicyl acid or any other components of this medicinal product stated in section 6
- On contact sensitisation,
- On babies,
- In the case of limited renal function.

Warnings and safety precautions for use

Callus Ointment Salicylic Vaseline 10 % should not be used on mucus membranes or near the eyes.

Using Callus Ointment Salicylic Vaseline 10 % with other medication

Please inform your doctor or pharmacist if you use, or have recently used, or intend to use other medicinal products. Salicylic acid can increase the absorption of other medicinal products applied on the same area. When absorbed by the body, salicylic acid delays the breakdown of methotrexate and increases the effect of sulfonylurea.

Pregnancy, lactation and fertility

Please ask your doctor or pharmacist for advice before use if you are pregnant or are nursing, if you think you might be pregnant or intend to become pregnant.

Do not use during pregnancy. Do not apply to the breast area during lactation. When treating corns, only use on areas up to 5 cm².

3. How should Callus Ointment Salicylic Vaseline 10 % be used?

Always use this medicinal product exactly as described in this leaflet or exactly as prescribed by your doctor, pharmacist or medical specialist. Please consult your doctor, pharmacist or medical specialist if you are unsure.



The recommended dose is: Apply to the skin 1-2 x daily.

Maximum daily dose for adults: less than 20 g (less than 2 g active ingredient).

Maximum daily dose for adolescents from 12 years of age: maximum 2 g (less than 0.2 g active ingredient).

How to use: For application to the skin.

Duration of use: Do not use for longer than one week.

Please consult your doctor or pharmacist or medical specialist if you have any further questions on taking this medicinal product.

4. Which side effects are possible?

As is the case with all medicinal products, this medicinal product can have side effects, but they do not necessarily affect everyone.

Skin irritation may occur on the areas where callus ointment has been applied.

In very rare cases, a previously unknown hypersensitivity (contact sensitisation) to salicylic acid may occur.

Reporting side effects

Please inform your doctor or pharmacist or medical specialist if you notice side effects. This also applies to side effects not stated in this leaflet. You can register any side effects directly with the Bundesinstitut für Arzneimittel und Medizinprodukte, Abt. Pharmakovigilanz, Kurt-Georg-Kiesinger-Allee 3, D-53175 Bonn, website: www.bfarm.de. By reporting any side effects, you can make a contribution towards more information being available on the safety of this medicinal product.

5. How should Callus Ointment Salicylic Vaseline 10 % be stored?

Keep this medicine out of the reach of children.

Do not take this medicinal product if the "use by" date on the carton of the container has expired. The use by date relates to the last day of the month in question.

Storage conditions: Do not store above 25 °C.

Never dispose of medicinal products in your waste water (e.g. not down the toilet or washbasin). Ask your pharmacist how you should dispose of the medicinal product if you no longer use it. This helps you contribute towards protection of the environment. More information is available at www.bfarm.de/arzneimittelentsorgung.

6. Pack contents and additional information

1 g ointment contains: Active ingredient: 0.1 g salicylic acid.

Other ingredients: Thin paraffin, white vaseline.

Packet sizes available: Callus Ointment Salicylic Vaseline 10 % is available with 90 g in the original pack.

This information leaflet was last updated in July 2021.

PB0222

Art. No. 894

For risks and side effects please read the information leaflet and consult your doctor or pharmacist.